

Elementary Background in Elliptic Curves

Antonia W. Blucher

National Security Agency, 9800 Savage Road, Fort George G. Meade, MD 20755-6000

Abstract. This paper gives additional background in algebraic geometry as an accompaniment to the article, “Formal Groups, Elliptic Curves, and some Theorems of Couveignes”. Section 1 discusses the addition law on elliptic curves, and Sections 2 and 3 explain about function fields, uniformizers, and powerseries expansions with respect to a uniformizer.

1 Elliptic curves

If K is a field then the 2-dimensional projective space $\mathbf{P}^2(K)$ is the set of triples $(X, Y, Z) \in K \times K \times K$ such that X, Y, Z are not all zero, modulo the equivalence $(X, Y, Z) \cong (lX, lY, lZ)$ for all $0 \neq l \in K$. Three points (X_i, Y_i, Z_i) ($i = 1, 2, 3$) in $\mathbf{P}^2(K)$ are said to be **colinear** if there exist $\lambda, \mu, \nu \in K$, not all zero, such that $lX_i + \mu Y_i + \nu Z_i = 0$ for $i = 1, 2, 3$. In particular, if $Z_i \neq 0$ for all i then the colinearity of the three points (X_i, Y_i, Z_i) in $\mathbf{P}^2(K)$ is equivalent to the colinearity of the points (x_i, y_i) in the affine plane, where $x_i = X_i/Z_i$ and $y_i = Y_i/Z_i$. This follows from the equation $lx_i + \mu y_i + \nu = 0$ for all i .

A **Weierstrass equation** is a cubic equation $W(X, Y, Z) = 0$, where

$$W(X, Y, Z) = Y^2Z + a_1XYZ + a_3YZ^2 - (X^3 + a_2X^2Z + a_4XZ^2 + a_6Z^3), \quad (1.1)$$

$a_i \in K$. Since this equation is homogeneous, we can think of its solutions as lying in $\mathbf{P}^2(K)$. A **singular point** is a point $P \in \mathbf{P}^2(\overline{K})$ such that $W(P) = \partial W/\partial X(P) = \partial W/\partial Y(P) = \partial W/\partial Z(P) = 0$, where $\overline{K}$ denotes the algebraic closure of K . It can be shown that the Weierstrass equation has no singular points iff $\Delta \neq 0$, where Δ is a certain polynomial in the a_i ; see [3], p. 46. Let us assume $\Delta \neq 0$. Then the set of solutions to (1.1) in $\mathbf{P}^2(\overline{K})$ is called an **elliptic curve**. The elements of the elliptic curve are called **points**. If L is any field containing all the Weierstrass coefficients a_i then $E(L)$ is defined to be the set of solutions to (1.1) in $\mathbf{P}^2(L)$. The Weierstrass equation can often be simplified. For example, in characteristic zero a linear change of coordinates puts the Weierstrass equation into the form $Y^2Z = X^3 + a_4XZ^2 + a_6Z^3$. See [3], Appendix A.

The only point on the elliptic curve which intersects the line $Z = 0$ is the point $(0, 1, 0)$. This is called the **point at infinity** and is often denoted by O_E . In contrast, the “finite” points of $E(K)$ are the points $(X, Y, 1)$ which satisfy the above equation, and it is customary to identify these points with the points (X, Y) in the affine plane $K \times K$ such that

$$Y^2 + a_1XY + a_3Y = X^3 + a_2X^2 + a_4X + a_6. \quad (1.2)$$

$E(K)$ can be made into an abelian group with additive identity equal to O_E by insisting that any three colinear points on $E(K)$ (counting multiplicities) sum to O_E . The proof that this group law is associative can be found in [2] using Bézout's Theorem or in [3] using the Picard group. In this context, the phrase “counting multiplicities” means to count the multiplicities of the roots of the cubic which results when one variable is eliminated from the Weierstrass equation by using the equation of the intersecting line. We give several examples which should make this clear.

Example 1.1: Lines through the origin. The equation of a line through $(0, 1, 0)$ has the form $\lambda X + \nu Z = 0$, where $\lambda, \nu \in K$ are not both zero. If $\lambda = 0$ then the line is $Z = 0$, and the Weierstrass equation becomes simply $0 = W(X, Y, 0) = X^3$. This has one solution $X = 0$ with multiplicity three. Thus $(0, 1, 0)$ intersects $Z = 0$ with multiplicity three. This gives $3(0, 1, 0) = (0, 1, 0)$, which is consistent with the fact that $(0, 1, 0)$ is the identity. If $\lambda \neq 0$ then $X = cZ$, where $c = -\nu/\lambda$. The Weierstrass equation becomes

$$W(cZ, Y, Z) = Y^2Z + (a_1c + a_3)YZ^2 - (c^3 + a_2c^2 + a_4c + a_6)Z^3 = 0.$$

This factors as

$$Z(Y^2 + (a_1c + a_3)YZ - (c^3 + a_2c^2 + a_4c + a_6)Z^2) = Z(Y - y_1Z)(Y - y_2Z)$$

where y_1, y_2 belong to a quadratic extension of K and satisfy $y_1 + y_2 = -(a_1c + a_3)$, $y_1y_2 = -(c^3 + a_2c^2 + a_4c + a_6)$. The intersection of E with the line is $\{(0, 1, 0), (c, y_1, 1), (c, y_2, 1)\}$, hence $(0, 1, 0) + (c, y_1, 1) + (c, y_2, 1) = (0, 1, 0)$. In other words, if $(c, y_1, 1)$ lies on E then $-(c, y_1, 1) = (c, y_2, 1)$, where $y_1 + y_2 = -(a_1c + a_3)$. This gives the inversion formula:

$$-(x, y, 1) = (x, -(y + a_1x + a_3), 1). \quad (1.3)$$

If it happens that $y = -(y + a_1x + a_3)$, that is, $2y = -(a_1x + a_3)$, then $(x, y, 1)$ is a two-torsion point.

Example 1.2: Lines which miss the origin. The equation of a line which misses $(0, 1, 0)$ is $\lambda X + \mu Y + \nu Z = 0$, where $\mu \neq 0$. Since $(0, 1, 0)$ is the only point on E which intersects $Z = 0$, the intersection points of the line with the elliptic curve all have a nonzero Z -coordinate. Let $m = -\lambda/\mu$, $b = -\nu/\mu$, $y = Y/Z$, $x = X/Z$. Then the three intersection points $(x_i, y_i, 1)$, $i = 1, 2, 3$, satisfy the Weierstrass equation and $y_i = mx_i + b$. Another way to view this is: suppose $(x_1, y_1, 1)$ and $(x_2, y_2, 1)$ are two points on the elliptic curve such that $x_1 \neq x_2$, and define $m = (y_1 - y_2)/(x_1 - x_2)$, $b = y_1 - mx_1$. Substitute $y = mx + b$ into the affine form of the Weierstrass equation to get a cubic equation in x of the form $x^3 + Bx^2 + Cx + D = 0$, where the coefficients explicitly depend on the a_i and on m and b . Explicitly, $B = -m^2 - a_1m + a_2$. Two roots of this cubic equation are x_1 and x_2 . Let x_3 denote the third root. Then

$$x^3 + Bx^2 + Cx + D = (x - x_1)(x - x_2)(x - x_3) = x^3 - (x_1 + x_2 + x_3)x^2 + (\cdots)x + (\cdots),$$

so that $x_3 = -x_1 - x_2 - B$. This yields the addition formula

$$(x_1, y_1, 1) + (x_2, y_2, 1) = (x_3, -(y_3 + a_1x_3 + a_3), 1) \quad \text{if } x_1 \neq x_2,$$

$$x_3 = -x_1 - x_2 + m^2 + a_1m - a_2, \quad m = (y_1 - y_2)/(x_1 - x_2),$$

$$y_3 = mx_3 + y_1 - mx_1.$$

Example 1.3: Duplication formula. How do we compute $P + P$? Assume $P = (x_1, y_1, 1)$ is not a two-torsion point, that is, $2y_1 \neq -(a_1x_1 + a_3)$. Then $-(P + P)$ is the point $(x_2, y_2, 1)$ on the curve such that $2(x_1, y_1, 1) + (x_2, y_2, 1) = (0, 1, 0)$. Let $lX + \mu Y + \nu Z$ be the line which passes through these points with correct multiplicity. It misses the origin, so $\mu \neq 0$. Thus the line has the form $y - y_1 = m(x - x_1)$, where $y = Y/Z$, $x = X/Z$. First we will find m , then we will find x_2, y_2 . When y is replaced by $m(x - x_1) + y_1$ in the Weierstrass equation, we get a cubic $x^3 + Bx^2 + Cx + D$, and it should equal $(x - x_1)^2(x - x_2)$. Here $B = a_2 - m^2 - a_1m$, $C = m^2(2x_1) + m(-2y_1 + a_1x_1 - a_3) + (a_4 - a_1y_1)$. If we differentiate the cubic and set $x = x_1$ we are supposed to get zero. Hence $3x_1^2 + 2Bx_1 + C = 0$. We can solve for m in this relation; in fact m will just be the slope of the tangent line to the curve E at P , namely $m = (dy/dx)(P)$. It turns out

$$m = (3x_1^2 + 2a_2x_1 + a_4 - a_1y_1)/(a_1x_1 + 2y_1 + a_3).$$

Since $(x - x_1)^2(x - x_2) = x^3 + Bx^2 + Cx + D$,

$$x_2 = -2x_1 - B = -2x_1 - a_2 + m^2 + a_1m, \quad y_2 = m(x_2 - x_1) + y_1.$$

Finally,

$$2(x_1, y_1, 1) = -(x_2, y_2, 1) = (x_2, -(y_2 + a_1x_2 + a_3), 1).$$

2 Function fields, local rings, and uniformizers

Let K be any field, and let $C = C(X, Y, Z)$ be a curve in $\mathbf{P}^2(\overline{K})$. This means that C is an irreducible homogeneous polynomial with coordinates in $\overline{K}$. Assume the coefficients of C belong to K , and let $C(K) = \{P \in \mathbf{P}^2(K) \mid C(P) = 0\}$. A point in $C(\overline{K})$ is called **nonsingular** if $\nabla_P C \neq 0$, where

$$\nabla_P C = \left(\frac{\partial}{\partial X} C(P), \frac{\partial}{\partial Y} C(P), \frac{\partial}{\partial Z} C(P) \right).$$

The **function field** $K(C)$ of C over K is the set of all quotients

$$p(X, Y, Z)/q(X, Y, Z)$$

, where p and q are homogeneous polynomials of the same degree with coefficients in K such that C does not divide q , modulo the equivalence: $p/q \cong p'/q'$ iff

$pq' - qp'$ is divisible by C . The addition and multiplication in $K(C)$ are defined in the same way as for rational functions. Equivalently, if we set $x = X/Z$ and $y = Y/Z$ then $K(C)$ is the quotient field of the integral domain

$$K[x, y]/(C(x, y, 1)).$$

This field has transcendence degree 1, thus any two elements of the function field satisfy some algebraic relation. Notice that $p(lX, lY, lZ)/q(lX, lY, lZ) = p(X, Y, Z)/q(X, Y, Z)$ because of the homogeneity, so $p(P)/q(P)$ makes sense for $P \in C(\overline{K})$ provided $q(P) \neq 0$. A function $f \in K(C)$ is said to be **defined** at a point $P \in C(\overline{K})$ if there exists p/q in the equivalence class of f such that $q(P) \neq 0$. If f is defined at P then the **value** of f at P , denoted $f(P)$, is defined as $p(P)/q(P)$ for any (hence every) p/q which is in the equivalence class of f and for which $q(P) \neq 0$. If $P \in C(K)$ and $f \in K(C)$ is defined at P then $f(P) \in K$.

As an example, let us show that for the Weierstrass equation, Z/X is defined at $(0, 1, 0)$. Since $Z(Y^2 + a_3YZ - a_6Z^2) = X(-a_1YZ + X^2 + a_2XZ + a_4Z^2)$,

$$Z/X = \frac{-a_1YZ + X^2 + a_2XZ + a_4Z^2}{Y^2 + a_3YZ - a_6Z^2}.$$

Now $Y^2 + a_3YZ - a_6Z^2$ does not vanish at $(0, 1, 0)$, so Z/X is indeed defined at $(0, 1, 0)$.

For the remainder of this section assume $P \in C(K)$. Let Ω_P be the ring of functions in $K(C)$ which are defined at P , and let M_P be the ideal of Ω_P consisting of functions which take the value zero at P . Ω_P is called the **local ring of C at P** . If we identify the set of constant functions with K then

$$\Omega_P = K \oplus M_P \tag{3.1}$$

(internal direct sum) because $f = f(P) + (f - f(P))$.

Lemma 2.1 $\Omega_P - M_P = \Omega_P^\times$. M_P is the unique maximal ideal of Ω_P .

Proof. If $f \in \Omega_P - M_P$ then f can be written as $F(X, Y, Z)/G(X, Y, Z)$, where F, G are homogeneous polynomials of the same degree and F, G do not vanish at P . Since $1/f$ can be written G/F , $1/f \in \Omega_P$. This shows that $\Omega_P - M_P \subset \Omega_P^\times$. The reverse inclusion is also true, for if $f \in \Omega_P^\times$ then there is $g \in \Omega_P$ with $fg = 1$. Then $f(P)g(P) = 1$, so $f(P)$ is a unit in K . In particular, f is not in M_P . We have proved the first statement. For the second statement, let I be any proper ideal of Ω_P . Then $I \cap \Omega_P^\times = \emptyset$; otherwise I would equal Ω_P . Thus I misses the complement of M_P completely; equivalently, $I \subset M_P$. $\square$

Our next goal is to show that if P is a nonsingular point (that is, $\nabla_P C \neq 0$) then M_P is principal. First we need some lemmas. These lemmas are true in more generality than we state them.

Lemma 2.2 (Nakayama's Lemma) If A is a finitely generated Ω_P -module and $M_P A = A$ then $A = \{0\}$.

Proof. Let $u_1, \dots, u_n$ be a set of generators for A with n as small as possible. Since $u_1 \in A = M_P A$, there exist $\mu_i \in M_P$ such that $u_1 = \sum \mu_i u_i$. Now $1 - \mu_1 \in \Omega_P^\times$, so we can solve for u_1 in terms of $u_2, \dots, u_n$. This contradicts the minimality of n . $\square$

Corollary 2.3 $\cap_{n=1}^\infty M_P^n = \{0\}$. Thus $M_P \supsetneq M_P^2$.

Proof. Let $A = \cap M_P^n$. A is finitely generated by the Hilbert Basis Theorem [1], p. 13, so $A = \{0\}$ by Nakayama's Lemma. $\square$

Lemma 2.4 If P is a nonsingular point of C then M_P/M_P^2 is isomorphic to K .

Proof. Observe that M_P/M_P^2 is an Ω_P/M_P -vector space, and Ω_P/M_P is canonically isomorphic to K by (3.1). We just have to prove this vector space is one-dimensional, or equivalently that its dual is one-dimensional. So let $l : M_P/M_P^2 \rightarrow K$ be a K -linear map. Since $\Omega_P = K + M_P$, we can think of l as a K -linear map from Ω_P into K which is trivial on $M_P^2 + K$.

We will construct a K -linear map $\vartheta : \Omega_P \rightarrow K$ which is trivial on $K + M_P^2$, and we will prove that λ is proportional to ϑ . Fix coordinates for P : $P = (X_0, Y_0, Z_0)$. Fix $R \in K^3$ such that $R \cdot P = 1$. Let $C(X, Y, Z) = 0$ be the equation of the curve. Let $\nabla_P C = (\partial C/\partial X(P), \partial C/\partial Y(P), \partial C/\partial Z(P))$, the gradient of C at P . We claim $\nabla_P C$ and R are linearly independent. First observe $\nabla_P C \neq 0$ since the curve is nonsingular at P by hypothesis. By Euler's identity $(X\partial/\partial X + Y\partial/\partial Y + Z\partial/\partial Z)C = \deg(C)C$. (The proof of Euler's identity is that for any monomial $X^a Y^b Z^c$, we have $(X\partial/\partial X + Y\partial/\partial Y + Z\partial/\partial Z)(X^a Y^b Z^c) = (a + b + c)X^a Y^b Z^c$.) Evaluating Euler's identity at P gives $P \cdot \nabla_P C = \deg(C)C(P) = 0$. Since $P \cdot R \neq 0$, this shows $\nabla_P C$ is linearly independent from R . It follows that the space of vectors which are orthogonal to R and $\nabla_P C$ is one-dimensional, spanned by a vector T .

Define $\vartheta : \Omega_P \rightarrow K$ by

$$\vartheta(f) = T \cdot \nabla_P(f).$$

We claim ϑ is well-defined, K -linear, and surjective. To prove it is well-defined, it suffices to show $T \cdot \nabla_P(F/G) = 0$ if F vanishes identically on the curve and $G(P) \neq 0$. In that case, C divides F . Let $h = F/(CG)$. Then

$$T \cdot \nabla_P(F/G) = C(P)T \cdot \nabla_P h + h(P)T \cdot \nabla_P C = 0 + 0$$

as required. The map ϑ is certainly K -linear. If S is any vector in K^3 such that $S \cdot P = 0$ but $S \cdot T \neq 0$ then

$$T \cdot \nabla_P(S \cdot (X, Y, Z)/R \cdot (X, Y, Z)) = \frac{T \cdot S}{R \cdot P} - \frac{(T \cdot R)(S \cdot P)}{(R \cdot P)^2} = \frac{T \cdot S}{R \cdot P} \neq 0, \quad (3.1)$$

so the map is nontrivial. Any nontrivial K -linear map into K is surjective. Let us show that the kernel contains $K + M_P^2$. Certainly $T \cdot \nabla_P$ annihilates the constants K . If $f, g \in M_P$ then $\nabla_P(fg) = 0$ by the product rule, so M_P^2 is in the kernel also.

It remains to show that an arbitrary linear map $l : \Omega_P \rightarrow K$ which is trivial on $K + M_P^2$ coincides with a multiple of ϑ . The map λ obeys a product rule, for if $f = f(P) + f_1$, $g = g(P) + g_1$ where $f, g \in \Omega_P$ then $f_1, g_1 \in M_P$, and $fg = f(P)g + g(P)f - f(P)g(P) + f_1g_1$, so

$$l(fg) = f(P)l(g) + g(P)l(f).$$

It follows that l is completely determined by its values at X/ρ , Y/ρ , and Z/ρ , where $\rho = R \cdot (X, Y, Z)$. Denote these by values by α, β, γ . We claim (α, β, γ) is proportional to T . This will imply there is a one-dimensional space of such λ ; in particular l must be proportional to ϑ . We must show (α, β, γ) is orthogonal to both R and $\nabla_P C$. Let $R = (R_1, R_2, R_3)$ and $\rho = R_1X + R_2Y + R_3Z$. Then

$$\begin{aligned} R \cdot (\alpha, \beta, \gamma) &= R_1l(X/\rho) + R_2l(Y/\rho) + R_3l(Z/\rho) \\ &= l((R_1X + R_2Y + R_3Z)/\rho) = l(1) = 0, \end{aligned}$$

$$\nabla_P C \cdot (\alpha, \beta, \gamma) = l(C(X/\rho, Y/\rho, Z/\rho)) = l(C(X, Y, Z)/\rho^{\deg(C)}) = 0.$$

□

Corollary 2.5 *Suppose P is nonsingular. Let $u \in M_P$, and let R, T be nonzero vectors in K^3 such that $R \cdot P \neq 0$, $T \cdot R = 0$, and $T \cdot \nabla_P C = 0$. Then $u \in M_P - M_P^2$ iff $T \cdot \nabla_P u \neq 0$. Also, if $S \cdot P = 0$ but $S \cdot T \neq 0$ then $S \cdot (X, Y, Z)/R \cdot (X, Y, Z)$ belongs to $M_P - M_P^2$.*

Proof. In the notation of the preceding proof, the hypothesis states that $\vartheta(u) \neq 0$. Since ϑ induces an isomorphism of M_P/M_P^2 with K , we know $u \notin M_P^2$ iff $\vartheta(u) \neq 0$. The last statement follows from (3.1). □

Proposition 2.6 *Suppose P is nonsingular. Then M_P is a principal ideal of Ω_P . In fact, if $u \in M_P$ then $M_P = u\Omega_P \iff u \notin M_P^2$.*

Proof. $M_P \neq M_P^2$ by Lemma 2.2. Let $u \in M_P - M_P^2$. Then

$$K \cong \frac{\Omega_P}{M_P} \cong \frac{u\Omega_P}{uM_P} \rightarrow \frac{u\Omega_P}{M_P^2}.$$

The image of 1 in this composite map is $u \bmod M_P^2$, which is nonzero. Thus the kernel of the projection map from $u\Omega_P/uM_P$ into $u\Omega_P/M_P^2$ has to be trivial. This shows $uM_P = M_P^2$.

Also, $M_P = uK + M_P^2$ because M_P/M_P^2 is a one-dimensional K -vector space by Lemma 3.4. Thus

$$u\Omega_P = u(K + M_P) = uK + uM_P = uK + M_P^2 = M_P.$$

□

Corollary 2.7 Suppose P is nonsingular. Let $0 \neq f \in K(C)$, $u \in M_P - M_P^2$. There is a unique integer $v(f)$ such that $f = u^{v(f)}g$ with $g \in \Omega_P^\times$. This integer does not depend on the choice of u .

Proof. Write $f = F/G$ with F, G homogeneous of degree m . Choose $R \in K^3$ so that $P \cdot R \neq 0$. Let ρ be the homogeneous polynomial $\rho = R \cdot (X, Y, Z)$. Then $f = f_1/f_2$, where $f_1 = F/\rho^m$ and $f_2 = G/\rho^m$. Note that $f_1, f_2 \in \Omega_P$. By Corollary 3.3,

$$\Omega_P - \{0\} = \sqcup_{n=0}^{\infty} (M_P^n - M_P^{n+1}),$$

where $M_P^0 = \Omega_P$. Clearly $M_P^n - M_P^{n+1} = u^n \Omega_P^\times$, since $M = u\Omega_P$. Define n_1, n_2 by $f_i \in M_P^{n_i} - M_P^{n_i+1}$, and let $n = n_1 - n_2$. Then $fu^{-n} = f_1u^{-n_1}/(f_2u^{-n_2}) \in \Omega_P^\times$. $\square$

Ω_P is called the **local ring** at P . If P is nonsingular, then a **uniformizer** is an element of M_P such that $M_P = u\Omega_P$; equivalently it is an element of $M_P - M_P^2$. The integer $v(f)$ is called the **valuation** of f at P . If $v(f) > 0$, f has a **zero of order $v(f)$ at P** ; if $v(f) < 0$, f has a **pole of order $-v(f)$ at P** . If $v(f) = 0$, we say f is **finite and nonvanishing at P** .

Example 2.8 Let $C = W$ be given by the Weierstrass equation (1.1). Let us show that X/Y is a uniformizer at $(0, 1, 0)$. In Corollary 2.5 we take $R = P = (0, 1, 0)$ and compute from (1.1) that $\nabla_P W = (0, 0, 1)$. Thus $T = (1, 0, 0)$. Now X/Y vanishes at P , and $T \cdot \nabla_P(X/Y) = \partial/\partial X(X/Y)|_P = 1 \neq 0$. So X/Y is a uniformizer by Corollary 2.5. $\square$

Example 2.9 Let us compute $v(Z/X)$ at the point $P = (0, 1, 0)$ on an elliptic curve. By (1.1), $ZV = X^3$, with

$$V = Y^2 + a_1XY + a_3YZ - a_2X^2 - a_4XZ - a_6Z^2.$$

Then $Z/X = X^2/V = (X/Y)^2(Y^2/V)$. Since $Y^2/V \in \Omega_P^\times$ and X/Y is a uniformizer, we see that $v(Z/X) = 2$. Thus $x = X/Z$ has a pole of order 2 at the origin. Since $Y/Z = (Y/X)(X/Z)$, $y = Y/Z$ has a pole of order 3 at the origin. $\square$

3 Power series expansions

Lemma 3.1 Let $C = C(X, Y, Z)$ be a projective curve over $\overline{K}$ which is nonsingular at P . Suppose C has coefficients in K and $P \in C(K)$. Let $u \in K(C)$ be a uniformizer at P . If f is a nonzero function in $K(C)$ and $v(f) = a$ then there exist unique constants $f_j \in K$ for $j \geq a$ such that for any $N \geq a$,

$$f - \sum_{j=a}^N f_j u^j \quad \text{has a zero of order at least } N + 1.$$

Moreover, $f_a \neq 0$.

Proof. Let $g = u^{-a}f \in \Omega_P^\times$, $f_a = g(P)$. This is the only constant for which $g - f_a \in M_P$, so it is the only constant for which $v(f - u^a f_a) = v(u^a(g - f_a)) > a$. From this observation, the lemma may be easily proved by induction. $\square$

Corollary 3.2 *A choice of uniformizer u at P determines a one to one homomorphism of rings*

$$\Psi_u : \Omega_P \hookrightarrow K[[\tau]].$$

If $0 \neq V = \sum v_i \tau^i \in K[[\tau]]$, define $\deg(V)$ to be the smallest integer n such that $v_n \neq 0$, and define $\deg(0) = \infty$. If one puts metrics on Ω_P and on $K[[\tau]]$ by the rules $|f| = c^{v_P(f)}$ for $f \in \Omega_P$, $|V| = c^{\deg(V)}$ for $V \in K[[\tau]]$, where $0 < c < 1$, then Ψ_u is an isometry. In particular, Ψ_u is continuous with respect to the topologies on Ω_P , $K[[\tau]]$ which are induced by the above metrics. The image of Ψ_u is dense in $K[[\tau]]$.

Proof. The homomorphism Ψ_u is defined by: $\Psi_u(f) = \sum f_j \tau^j$, where the f_j are as in Lemma 3.1. It is easy to see (using the uniqueness of the coefficients f_j) that Ψ_u is a ring homomorphism. Ψ_u is an isometry because $v_P(f) = a$ implies f_a is the first nonzero coefficient. If $\Psi_u(f) = 0$ then $|f| = |\Psi_u(f)| = |0| = 0$, so $f \in \cap M_P^n$. By Corollary 2.3, f must be zero; thus Ψ_u is one to one. If g is any polynomial with coefficients in K then $g(u) \in \Omega_P$ and $\Psi_u(g(u)) = g(\tau)$. This shows $\Psi_u(\Omega_P)$ is dense in $K[[\tau]]$. $\square$

References

1. W. Fulton, *Algebraic Curves*, Addison-Wesley, 1969
2. A. W. Knap, *Elliptic Curves*, Math. Notes **40**, Princeton University Press, Princeton, New Jersey, 1992
2. J. H. Silverman, *The Arithmetic of Elliptic Curves*, Springer-Verlag, New York, 1986